



Age structure of Patagonian toothfish in the Falkland Islands

2014 - 2015

Brendon Lee

Directorate of
Natural Resources
Fisheries



TABLE OF CONTENTS

1. Introduction	1
2. Methods	1
2.1. Data Collection.....	1
2.2. Preparation of otoliths.....	2
2.3. Reading methodology	2
2.4. Estimation of von Bertalanffy parameters	2
2.5. Precision of the age estimates.....	3
3. Results and discussion.....	4
3.1. Distribution of Samples.....	4
3.2. Length and Age composition.....	5
3.3. Size and Age	6
3.4. Precision of the age estimates.....	8
4. References	10
5. Acknowledgements.....	10
6. Appendix A	11

AGE STRUCTURE OF TOOTHFISH - 2014

1. Introduction

Growth parameters, estimates of age at maturity and recruitment, and population age structures are essential inputs in fisheries yield models that provide the basis for management advice in many world fisheries (Ashford, 2002, Horn, 2002).

Methodology for age determination should be applied across the catch and over a sequence of years while minimizing sources of error (Morison et al., 1998). In addition, quantitative estimates of variability in age determination should be provided for incorporation into decisions on analyses and modelling and interpretation.

This annual report, therefore presents a reliable ageing methodology for Patagonian toothfish, *Dissostichus eleginoides*, to calculate growth parameters from samples obtained from both trawl caught pre-recruits and from adult fish taken in the longline fishery in the Falkland Islands during 2014. It also presented an assessment of the bias and precision in order to ensure error does not exceed a quality threshold.

2. Methods

2.1. Data Collection

D. eleginoides were sampled by scientific observers and other scientific staff of the Falkland Islands Government Fisheries Department. Data were collected on board licensed longliners as well as on board commercial trawlers operating bottom trawls under various license types. In addition data were collected on board RV 'Castello' operating bottom and semi-pelagic trawls during two research cruises.

Randomly sampled toothfish were measured to the nearest cm (Lt), sexed and the stage of reproductive maturity assigned according to an eight-stage scale (I and II – immature, III and IV – maturing, V – mature, VI – running, VII – post spawning and VIII – spent). Otoliths were removed from a sub-sample of fish selected in order to cover the length range of the captured fish from within the various fisheries. Each annual collection of otoliths are stored in paper envelopes in four quarterly time periods (A: Jan – Mar, B: Apr – Jun, C: Jul – Sep and D: Oct – Dec).

AGE STRUCTURE OF TOOTHFISH - 2014

Otoliths for ageing are selected to cover the length distribution of sampled fish from each quarterly otolith collection. This ensures that sufficient otoliths are aged for all lengths on a temporal basis.

2.2. Preparation of otoliths

Otoliths were embedded in rows of five in blocks of amber coloured polyester resin and left to set for 24 hours. Fully dried blocks are ground in order to provide smooth linear surfaces and the nucleus marked using a pencil. This is undertaken in order to guide the cutting angle and ensure that sections are cut precisely at right angles. Resin blocks were subsequently sectioned using a Buehler Isomet Low Speed Saw. Between two and six sections of 0.35mm were taken per resin block and mounted on microscope slides under coverslips with clear polyester resin.

2.3. Reading methodology

Sections were viewed under reflected light at 10 to 40 times magnification. All sections of each row of otoliths were inspected and the section closest to the primordium was used for subsequent ageing. Images were taken for the best section for each otolith and saved for image enhancement and assistance in ageing.

Following previous work on age estimation of this species, the sector from the primordium to the proximal edge of the section, on the ventral side of the sulcus was chosen as the area in which to count increments. However, for some preparation, increments formed on the dorsal side were at least as clear as those on the ventral side. A readability index of 1 – Easy, 2 – Medium and 3 – Difficult was assigned to each otolith. Each otolith was aged twice by the primary reader. All counts of annuli were made without prior knowledge of fish size, date of capture or previous age estimates.

2.4. Estimation of von Bertalanffy parameters

Von Bertalanffy growth parameters were estimated from the length-at-age data using non-linear least squares regression procedure using R software (R core Team, 2015). 95% confidence was calculated for the von Bertalanffy parameters along with upper and lower 95% prediction limits of the derived curve. The age structure of the total sample of *D. eleginoides* captured in 2014 was estimated by constructing an

AGE STRUCTURE OF TOOTHFISH - 2014

age-length key (ALK) using the FSA package in R (Ogle, 2016). The relationship between length and weight was estimated using linear regression analysis

2.5. Precision of the age estimates

Repeated readings of the same otoliths provide a measure of intra-reader or inter-reader variability. They do not validate the assigned ages but provide an indication of size of the error to be expected with a set of age estimates, due to variation in interpretation of an otolith. Beamish and Fournier (1987) have developed an index of average percent error (IAPE), which has become a common method for quantifying this variation. The IAPE is calculated as:

$$IAPE = \frac{100}{N} \sum_{j=1}^N \left[\frac{1}{R} \sum_{i=1}^R \frac{|X_{ij} - X_j|}{X_j} \right]$$

Where N is the number of fish aged, R is the number of times fish are aged, X_{ij} is the i th determination for the j th fish, and X_j is the average estimated age of the j th fish.

Chang (1982) suggested that precision should be measured by the mean coefficient of variation (ACV) which is defined as:

$$ACV = 100 * \frac{\sum_{j=1}^n \frac{S_j}{X_j}}{n}$$

where s_j is the standard deviation of the R age estimates for the j th fish.

An IAPE and ACV were calculated for all repeated readings undertaken by the primary reader. The distributions of the differences between repeat readings were also inspected as another indicator of ageing errors, and of any bias between readings. Precision of repeated age estimates was also examined using an age bias plot (Campana *et al.*, 1995).

AGE STRUCTURE OF TOOTHFISH - 2014

3. Results and discussion

3.1. Distribution of Samples

During 2014, biological information was obtained from a total of 7761 *D. eleginoides* samples. Of these samples 3161 were obtained from within the trawl fishery and 4600 from within the longline fishery. Age data were obtained from a subsample of 679 fish.

D. eleginoides were captured in trawls and longlines over the Falkland Islands shelf and slope between 95 and 2009 m depth (Figure 1). In the trawl fishery, fish occurred between 95 and 592 m (average 243.05 m). In the longline fishery, samples were obtained from between 710 and 2009 m with an average of fepth of 1341 m.




Figure 1: Positions from which *D. eleginoides* samples were obtained over the Falkland Islands shelf and slope.

AGE STRUCTURE OF TOOTHFISH - 2014

3.2. Length and Age composition

The length frequency distribution of *D. eleginoides* sampled during 2014 was bimodal with an average total length of 72.02cm (Figure 2A). The trawl and longline fisheries targeted different parts of the *D. eleginoides* stock. The average length of fish captured within the trawl fishery was 30.94cm (25cm mode) while within the longline fishery the average length was 100.26cm (100cm mode).

The age distribution of *D. eleginoides* captured during 2014 as estimated from the age length key was bimodally distributed (Figure 2B). The average age of the total sample of *D. eleginoides* was 7.39 years. However, the trawl fishery captured younger fish (mean 1.83) compared to the longline fishery (11.2 years)




Figure 2: Length (A) and age frequencies (B) estimated from the total (aged and unaged) sampled catch of *D. eleginoides* in the Falkland Islands during 2014. Ages for unaged fish were obtained through the construction of an age-length-key.

Results of the length-weight regression indicate a significant relationship between length and weight ($P < 0.001$) with low variability around the line ($R^2 = 0.998$, Table 1, Figure 3).

AGE STRUCTURE OF TOOTHFISH - 2014

Table 1: Parameter estimates for the length-weight regression for *D. eleginoides* sampled during 2014.

Coefficient	Estimate	Std error	t-value	P
Intercept	-2.36	0.0047	-506.8	<0.001
LogL	3.18	0.0027	1201.2	<0.001
R2	0.998			




Figure 3: The log₁₀ – log₁₀ transformed length-weight data with the best fit line superimposed (left) and the length-weight data with the back-transformed best-fit line superimposed for *D. eleginoides* sampled during 2014.

3.3. Size and Age

D. eleginoides ages ranged from 0 to 33 years though fish older than 21 years were not common (Figure 4). Calculated von Bertalanffy growth parameters and their 95% confidence intervals are presented in Table 2 and Figure 5.

AGE STRUCTURE OF TOOTHFISH - 2014




Figure 4: Length versus age with superimposed best-fit von Bertalanffy growth model and 95% confidence bands (dashed lines) for *D. eleginoides* sampled during 2014.

Table 2: Von Bertalanffy parameters (with 95% confidence intervals) for *D. eleginoides* sampled during 2014.

Parameters	Estimate	Std Error	LCI	UCI
Linf	152.59	2.84	147.29	158.92
K	0.094	0.004	0.086	0.1
t0	-0.732	0.0892	-0.92	-0.57




Figure 5: Histograms of parameter estimates for the von Bertalanffy growth function from bootstrapped samples for *D. eleginoides* sampled during 2014.

AGE STRUCTURE OF TOOTHFISH - 2014

3.4. Precision of the age estimates

The percentage agreement table indicates that multiple estimates of ages by the primary reader agreed for 62.74% of the fish, 27.54% differed by one year and 7.81% differed by two years (Table 3). Figure 6 shows a tendency for the second age estimate to be lower than the accepted age estimate, particularly for older age estimates. The APE was 4.55% and the ACV was 6.43% (Table 4). These represent a satisfactory level of precision but suggest that the otoliths of *D. eleginoides* are not easy to age.

Table 3: Percentage table of raw differences between multiple readings of *D. eleginoides* otoliths for 2014.

Years	-4	-3	-2	-1	0	1	2
Frequency (%)	0.29	1.62	6.48	14.29	62.74	13.25	1.33




Figure 6: Age-bias plot for comparing the mean estimated ages from multiple readings of *D. eleginoides* for 2014.

AGE STRUCTURE OF TOOTHFISH - 2014

Table 4: Estimates of average coefficient of variation and average percent error for multiple ageings of *D. eleginoides* otoliths for 2014

n	Valid n	R	ACV	APE	Percent agreement
1461	679	2	6.43	4.55	62.74

AGE STRUCTURE OF TOOTHFISH - 2014

4. References

Ashford, J., Jones, C. and Bobko, S. 2002. Length-at-age in juvenile Patagonian toothfish (*Dissostichus eleginoides*). CCAMLR Science 9: 1 – 10.

Beamish, R. and Fournier, D. A. 1981. A method for comparing the precision of a set of age determinations. Journal of the Fisheries Research Board of Canada 36: 1395 – 1400.

Campana, S. E., Annand, M. C. and McMillan, J. I. 1995. Graphical and statistical methods for determining the consistency of age determinations. Transactions of the American Fisheries Society 124: 131 – 138.

Chang, W. Y. B. 1982. A statistical method for evaluating the reproducibility of age determination. Canadian Journal of Fisheries and Aquatic Sciences 39: 1208 - 1210.

Horn, P. L. 2002. Age and growth of Patagonian toothfish (*Dissostichus eleginoides*) and Antarctic toothfish (*D. mawsoni*) in waters from the New Zealand subantarctic to the Ross Sea, Antarctica. Fisheries Research 56: 275 – 287.

Ogle, D.H. 2016. Introductory Fisheries Analyses with R. Chapman & Hall/CRC, Boca Raton, FL.


R Core Team (2015). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

5. Acknowledgements

Thank you to the Scientific Fisheries Observers of the Falkland Islands Government Fisheries Department for the collection of biological data.

AGE STRUCTURE OF TOOTHFISH - 2014

6. Appendix A



Bubble plot representation of the observed age-length key for *D. eleginoides* sampled during 2014. The area of each circle is proportional to the proportion of fish in a length interval that are a given age.